

**NITROGEN UTILIZATION OF VEGETABLES GROWN
UNDER PLASTIC GREENHOUSE IN ANKARA CONDITIONS
USING ^{15}N TECHNIQUE**

Halitligil M. B., Kışlal H., Şirin H., Şirin C., Kiliçaslan A.
Turkish Atomic Energy Authority, Ankara, Turkey

ABSTRACT

In order to find suitable varieties of tomato, pepper and cucumber for plastic greenhouse conditions in Ankara and eventually to identify the best N fertilizer rate greenhouse experiments were conducted for two years. Yazgı F₁ variety for tomato, Hızır F₁ variety for cucumber and Serademre 8 variety for pepper were chosen to be the suitable varieties to grow in the plastic greenhouse conditions in Ankara.

Five N treatments [$N_0 = 0$, $N_1 = 150$, $N_2 = 300$, and $N_3 = 450$ kg N/ha; also, soil N application treatment (N_{soil}) equivalent to the fertigation treatment of 300 kg N/ha was included for tomato and pepper, however N rates for cucumber was 131, 266 and 339 kg N/ha; N_{soil} being 266 kg N/ha.] were investigated using ^{15}N labeled urea fertilizer.

Significantly higher marketable fresh fruit and total dry matter yields and N uptakes values were obtained from N_3 treatments for tomato and cucumber, but from N_2 treatment for pepper. Also, significantly higher yields, N uptakes and % NUE values were obtained when the same amount of N fertilizer is applied through fertigation compared to the treatment where N fertilizer applied to the soil then drip irrigated.

INTRODUCTION

The plants can use nitrogen easily and more efficiently if they can exist at optimum levels in the effective root zone of the plants. Drip irrigation-fertigation is the system that can provide both nitrogen and water at desired levels and conditions in the root zone. Both yields, nitrogen and water use efficiencies can be increased by drip irrigation-fertigation [1,2,3]. This system had been as a sensible method of supplying nitrogen inside the greenhouse [2,3]. Plastic greenhouses without heating systems had been widely used in the Mediterranean and Aegean Sea Regions of Turkey successfully during the last decade for drip irrigated–fertigated vegetable production [3]. However, plastic greenhouses are recently had being used for vegetable production at temperate locations of Central Anatolia, like Ankara, without the need of any additional heating in cool weather (early April – late October). In general, open field vegetables are produced and sent to market in mid July in Ankara. To grow vegetables earlier than July in Ankara climatic conditions will be an extra income to the farmers due to selling their vegetables for higher prices.

The specific objective of this study was to compare the nitrogen fertilizer use efficiency under conventional N fertilizer application with N fertigation when drip irrigation system is used in the greenhouse conditions in Ankara.

MATERIAL AND METHODS

In order to find out the suitable varieties of tomato, pepper and cucumber for plastic greenhouse conditions in Ankara, four different varieties from each vegetable, namely, Tomato (Ecem F_1 , 9920 F_1 , 2116 F_1 and Yazgı F_1), Cucumber (Hızır F_1 , Rapido, Hana, and Luna) and Pepper (1245 F_1 , 730 F_1 , Serademre 8 and 710 F_1) varieties were grown in the plastic greenhouse of Ankara Nuclear Research Center for Agriculture and Animal Sciences (ANRCAAS) at Saray-Ankara in 2001. Due to the higher marketable fresh fruit and total dry matter yields, N uptakes and NUE values obtained from Yazgı F_1 , Hızır F_1 and Serademre 8 varieties for tomato, cucumber and pepper, respectively, were chosen to be used in 2002 and 2003 experiments.

The soil used in the experiment had a pH of 7.9 and was low in organic matter content. Greenhouse was divided into three sections and in each section there were two 70 cm wide ridges and in between the ridges there was one 70 cm furrow for each vegetable. For every vegetable section there were four rows and at each row plants were 70 cm apart which ended up with a plant density of 20 000 plants per hectare. Five N treatments were investigated in all experiments. Nitrogen as urea was applied through drip irrigation water at rates of $N_0 = 0$, $N_1 = 150$, $N_2 = 300$, and $N_3 = 450$ kg N/ha; also, soil N application treatment (N_{soil}) equivalent to the fertigation treatment of 300 kg N/ha was included for tomato and pepper, however N rates for cucumber was 131, 266 and 339 kg N/ha; N_{soil} being 266 kg N/ha.

Each vegetable experiment consisted of two blocks in which the five N treatments plots were established in a randomized complete block experimental design. Therefore, each N treatment plot consisted of two rows having 8 plants ($2.8 \text{ m} \times 1.4 \text{ m} = 3.92 \text{ m}^2$) at planting. The middle four plants were considered to be the replications and each of them was assigned with a

replication number. ^{15}N isotope subplots were established on each ridge including 8 plants with a different lateral dripper line. Urea fertilizer with % 1,61 ^{15}N atom excess enrichment was used.

At final harvest, each assigned replication plant from each plot was harvested separately and divided into fruit, leaf, stem and root parts, then dried at 70 °C, ground and passed 2 mm screen. Total N and ^{15}N analyses were done by using semi microKjeldahl and NOI7 emission spectrometer, respectively. Afterwards, Ndff and NUE values were calculated.

RESULTS AND DISCUSSION

Two year average values of yield, N uptake, nitrogen use efficiency (% NUE) for tomato variety Yazgı F₁, cucumber variety Hızır F₁ and pepper variety Serademre 8 at different N rates are given in Table 1.

Significantly higher marketable fresh fruit and total dry matter yields and N uptake values were obtained from N₃ treatments for tomato and cucumber, but from N₂ treatment for pepper. Significantly lower % NUE values were observed by decreasing the N rate, which is in accordance with the findings of other researchers [2,3]. Highest % NUE values were obtained for cucumber and the lowest values obtained for tomato. For all three vegetables the lowest marketable fresh fruit and total dry matter yields, N uptakes values were obtained when no N fertilizer applied (N₀ treatment).

Table 1. Two years average total yields, N uptakes, Ndff and % NUE of vegetables grown at different n treatments under plastic greenhouse conditions.

	N Rates				
	N ₀	N ₁	N ₂	N ₃	N _{soil}
TOMATO (Yazgı F₁) Fresh Fruit Yields (kg/ha)	146500 c	170000 ab	174930 a	181270 a	165630 b
Total Dry Matter Yields (kg/ha)	10003 c	11681 b	11692 b	12383 a	11402 b
Total N uptake (kg N/ha)	159 d	186 c	204 b	222 a	205 b
Ndff (kg N/ha)		44 d	79 b	110 a	73 c
% NUE		28.9 a	26.4 b	24.5 c	24.2 c
CUCUMBER (Hızır F₁) Fresh Fruit Yields (kg/ha)	107880 c	148100 b	157750 b	170400 a	154230 b
Total Dry Matter Yields (kg/ha)	7937 d	9797 c	10301 b	10535 a	10130 b
Total N uptake (kg N/ha)	158 d	202 c	242 a	250 a	222 b
Ndff (kg N/ha)		79 c	125 b	155 a	83 c
% NUE		59.8 a	47.6 b	39.3 c	31.7 d
PEPPER (Serademre 8) Fresh Fruit Yields (kg/ha)	72350 e	83100 d	111230 a	99250 b	89430 c
Total Dry Matter Yields (kg/da)	8155 d	10736 c	13175 a	12478 b	10860 c
Total N uptake (kg N/ha)	175 e	206 d	300 a	262 b	230 c
Ndff (kg N/ha)		47 d	106b	125 a	79 c
% NUE		30.5 b	36.5 a	28.8 bc	26.6 c

* Values followed by the same letter within a row are not significantly different at 0.05 levels according to LSD analysis.

Significantly higher yields, N uptakes, % NUE values were obtained when same amount of N fertilizer (N₂ treatment) is applied through fertigation compared to the treatment where N fertilizer applied to the soil then drip irrigated (N_{soil} treatment). In other words, nitrogen rate N₃ for tomato and cucumber, but nitrogen rate N₂ was superior in yields.

When two-year average % N and % Ndff (nitrogen derived from fertilizer) values obtained at harvest for different soil depths under different vegetables were evaluated, higher % N and % Ndff values were obtained as the N rate increased regardless of soil depth. Generally, lower % Ndff values are obtained as the soil depth increased and as it can be seen from the table no labeled N was detected in the soil samples taken from 60-80 cm depth. The % Ndff values

obtained for the conventional N-fertilizer application treatment (N_{soil}) were higher than the same amount of N-fertilizer applied through fertigation (N_2), which showed that with fertigation treatment less N fertilizer was leached.

CONCLUSION

With this study it was proven that without any heating, one month early (first pick in June) and one month late (last pick at late October) tomato and pepper production is feasible under plastic greenhouse conditions in Ankara. Although early production for cucumber is feasible in the plastic greenhouse late cucumber production is not feasible.

It was shown that Yazgı F₁, Hızır F₁ and Serademre 8 are the suitable varieties of tomato, cucumber and pepper, respectively, for the plastic greenhouse conditions in Ankara due to their higher yields and % NUE values. It was also shown that nitrogen fertilization is necessary for the higher productivity of tomato, cucumber and pepper and with control treatments (N_0) nearly 24-58 % lower marketable fresh fruit yields were obtained compared to the yields obtained from N_3 or N_2 treatments. Also, soil application of N fertilizer (N_{soil}) was less effective when compared to fertigation treatment with same N rate.

REFERENCES

1. Moutonnet, P. and Heng, L.K. (2002) Overview of the IAEA programme on fertigation studies in the Mediterranean region. In Nuclear techniques in Integrated Plant Nutrient, Water and Soil Management. Proceedings of an International Symposium, 16-20 October 2000. Vienna, Austria. IAEA-C&S 11 and IAEA-C&S-11/C. pp. 217-234.
2. Halitligil, M.B., Akın, A.I., Kışlal, H., Öztürk, A., Deviren, A. (2002) Yield, nitrogen uptake and nitrogen use efficiencies by tomato, pepper, cucumber, melon and eggplant as affected by nitrogen rates applied with drip-irrigation under greenhouse conditions. In Water Balance and Fertigation for Crop Improvement in West Asia. IAEA-TECDOC-1266, pp. 99-110.
3. Zuraiqi, S., Qawasmi, W., Deek, I., Mohammad, M.J. (1998) Management of nitrogen fertigation of tomato with the use of ^{15}N technology. In Water Balance and Fertigation for Crop Improvement in West Asia. IAEA-TECDOC-1266 Vienna (2002) pp. 27-40. Handbook. IAEA-Vienna 48 p.